

CABBAGE LARI



EARLY MATURING FRESH- MARKET HYBRID CABBAGE FOR AFRICAN CONDITIONS

- Very high yield potential
- Excellent uniformity
- Early maturing with good holdability
- Adaptable to summer conditions

Type

Early in maturity, medium size heads cabbage for fresh market.

Maturity

Early maturity. Approximately 75 - 85 days from transplant to maturity in summer.

Plant Characteristics

Strong plant vigour, medium frame size with a round head, has an open growth habit. Good uniformity in the field with a blue-green colour. Plant population of 30,000 to 35,000 plants per ha is recommended for single head marketing. Plant population will influence head size. For larger head size the plant population should be reduced to suit your needs.

Head Characteristics: LARI has very firm globe shaped heads with an average head size of between 3.5 – 4.5 kg with an excellent flavour. LARI has a very high first cut percentage and good standability.

Variety Characteristics

Fertilization Guidelines: This is a general guide, grower conditions vary and a soil analysis is highly recommended:

N: 150 to 200kg/ha. 70 to 90 kg/ha incorporated preplant.

Top dress the balance as follows:

Summer: 7,14 and 28 days after transplanting (no later).

Winter: 14,28 and 45 days from transplant (no later).

P: 50 kg/ha incorporated preplant. K: 120 kg/ha incorporate preplant. Soil pH: 6 - 6,8.

Features & Benefits

The combination of early maturity and vigorous homogenously emerging seedlings, contribute to the excellent uniform crop that LARI offers the grower.

Disease Resistance: Field observation shows good general disease resistance

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Starke Ayres Standard Terms and Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).