


TOMATO PEREGRINE



A DETERMINATE SALADETTE TOMATO WITH EXCELLENT FRUIT QUALITY ON A HIGHLY DISEASE RESISTANT PLANT

- Semi-determinate variety with high yield
- Extending bearing period
- Good disease resistance package
- Exceptional leaf disease tolerance
- Ideal for fresh and informal markets

Type:

A semi determinate processing tomato hybrid with an extended set suited to open-field production for the fresh and informal market as a trellised or untrellised type.

Maturity:

Medium maturing variety. In Summer plantings the first fruit will mature in 75 - 80 days after transplanting. The date of harvesting may vary by as much as 10 days between early and late season plantings. This needs to be taken into consideration when planning planting schedules under different growing conditions.

Plant Characteristics:

Vigorous with exceptional leaf cover on a highly disease tolerant plant with a semi- determinate growth habit.

Plant Population:

Peregrine could be grown without trellising, it is highly recommended to be trellised to 1 – 3 wires. The planting density should be chosen to give 15,000 - 21,000 plants per hectare. The row spacing should not be less than 35cm between plants. For an trellised production, a plant population of 10,000 - 15,000 plants per hectare with a row spacing of not less than 2 - 2.5m and an in-row spacing of 27 - 30 cm is recommended

Variety Characteristics:

Peregrine has a very high quality blocky/ cylindrical fruit with an average mass of 120 - 140 g. Fruit quality is exceptional with a deep red colour, thick walls, good flavour and no green shoulders clean shoulders. Peregrine have a very high marketable yield potential; a high percentage of large first grades can be expected.

Features & Benefits:

Peregrine is a Blye type hybrid with a very good disease resistance package and comparable yield that it maintains well later into the season. It also has very good size potential.

Disease Resistance:

HR: Verticillium Wilt (Va,Vd), Fusarium Wilt (Fol 1-3), Tobacco Mosaic Virus (TMV), Tomato Yellow Leaf Curl Virus (TYLCV).
IR: Root-knot Nematodes (Ma,Mi,Mj), Fusarium Crown & Root Rot (For), Tomato Spotted Wilt Virus (TSWV).

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Starke Ayres Standard Terms and Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).