

# CABBAGE RUBY KING



## QUICK-MATURING BIG RED CABBAGE FOR YEAR-ROUND PRODUCTION

- Round head with very good uniformity
- Quick-maturing: 65 to 75 days in summer
- Winter maturing from 80 to 100 days
- Good black rot resistance
- Year-round production



### Type

A big red hybrid cabbage variety for fresh and pre-pack market.

### Maturity

65 - 75 days (summer).  
85 - 100 days (winter).

### Plant Characteristics

RUBY KING has a medium-large frame size and good head size. Plant population will influence head size.

**Head Characteristics:** RUBY KING produces firm, round heads. Heads are uniform. Internal quality of the head is excellent.

### Variety Characteristics

**Fertilization Guidelines:** This is a general guide, grower conditions vary and a soil analysis is highly recommended.

N: 150 to 200kg/ha. 70 to 90 kg/ha incorporated preplant.

Top dress the balance as follows: 7,14 and 28 days after transplanting (no later).

P: 50 kg/ha incorporated preplant. K: 120 kg/ha incorporate preplant. Soil pH: 6 - 6,8.

### Features & Benefits

This variety shows good adaptability for year-round production in certain areas.

**Disease Resistance:** Field observation indicates some resistance to black rot.

**Disclaimer:** This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Starke Ayres Standard Terms and Conditions of Sale before ordering seed.

**Resistance:** is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).